

EFFECT OF SPROUTED FLAX SEEDS ON THE BODY'S REACTIVITY

Ismatova Marguba Shaukatovna

Assistant of the Department of Physiology of the Samarkand State Medical University

Rakhmatova Fotima Ulugbekovna

4th year student, group 431 Faculty of Medicine, Samarkand State Medical University

Samarkand, Uzbekistan

Annotation: This article examines the influence of sprouted flax seeds on the body's reactivity, biochemical composition, benefits and harms of flax seed. The percentage of fatty acids in flaxseed oil is compared with its content in other vegetable oils. The question of the possibility of using flax seeds for food and the content of cyanogenic glycosides in them is considered.

Key words: Flax seeds, fatty acids, amygdalin, cyanide, lignans.

Due to its unique properties, flax seeds attract close interest as a source of biologically active substances necessary for functional nutrition and maintaining of human health. They are rich in essential polyunsaturated fatty acids, dietary fibers, complete protein, polypeptides and lignans, belonging to the class of phytoestrogens that support the most important physiological functions of the human body.

It has always been believed that flax seed is one of the most useful products. They are rich in fiber, fatty acids and trace elements. Therefore, flax seeds are an integral part of proper nutrition - they are added to various salads and drinks, and many people eat them daily. The composition of flaxseed of Canadian varieties, which are the overwhelming majority in world production, is as follows: fat-41 %, protein-21 %, fiber-28 %, aromatic acids, sugars, lignin and hemicellulose-6 %, bottom ash-4 %. The composition of flaxseed varies depending on the variety, growing environment, and processing methods of flax.

A very significant part of flax seeds are lipids. Lipids are the components of cells that are difficult to dissolve in water. Fatty oil consists of triglycerides (natural organic compounds full of glycerol esters and monobasic fatty acids) and a mixture of fatty acids such as linoleic, linolenic, oleic, palmitic and stearic. The content of fatty acids in vegetable oils, % is given in Table 1.

Table № 1

Types of Oils	Saturated LC	Monounsaturated LC	Polyunsaturated LC	
		Oleic Acid (Omega-9)	Linoleic Acid (Omega-6)	Linolenic Acid (Omega-3)
Flaxseed	8-10	14	24-49	22-44
Soy	7,1-15,2	32,6-35,5	52-56	3-8

Olive	oil 9,0-14,3	71-86	4,2-11,8	-
Sunflower oil	9	33,3	39,8-60	-
Corn	oil 11,9	44,9-45,3	41-48	-
Hemp	oil 4,5	14	65	16

Plant fatty oils are a concentrated energy and building reserve, concentrated in seeds and other plant organs. The fat content in seeds and fruits of plants varies widely - from 2 to 70% and depends on the geographical location and climatic conditions in the area of their growth. The main role of reserve fats in the plant is to use them for nutrition during seed germination and embryo development; in addition, they play an important role as protective substances that help the plant to tolerate adverse environmental conditions, in particular, low temperatures. Unsaturated fatty acids have the greatest calorific value, so plants of northern latitudes contain them in the greatest quantities. In tropical plants, on the contrary, saturated fatty acids predominate, which are in a liquid state at high temperatures and solidify at low temperatures.

Flax seed lignans. Flaxseed is one of the richest sources of lignans belonging to the class of phytoestrogens. These are substances of plant origin that exhibit estrogen-like activity in the human body.

The main lignan of flaxseed – secoisolaracyresinol diglycoside (SDH) in the human body under the action of intestinal microflora metabolizes into enterolactone and enterodiol.

Numerous epidemiological studies have shown that flaxseed lignans reduce the risk of developing hormone-dependent tumors or slow their growth, and exhibit significant antioxidant activity. However, with regard to the use of flax seeds in food, you should not get carried away with them, as this can cause serious health problems.

Cyanide is a poison that acts quickly and can be fatal. There is a link between cyanide and flax. As it turned out, in addition to essential fatty acids, flax seed contains a compound called amygdalin, which is one of the "cyanogenic glycosides" that, when decomposed (when destroyed in the digestive tract), can produce cyanide gas. Most of it is contained in ground flaxseed.

Experts of the European Food Safety Agency (EFSA) have published a report in which they warn that the consumption of only 1.3 g (one-third of a teaspoon) of ground flax seed can pose a danger to the health of a small child, as the amount of cyanide in his body will reach dangerous levels. An adult may experience intoxication if they eat 10.9 g (three teaspoons) of flax seeds at a time. If the seed was crushed using a blender, mortar, or by hand, cyanide release occurs several times faster. At the same time, the more crushed the seed is, the more cyanide it emits.

Signs of cyanide poisoning are as follows: headache, absent-mindedness, agitation of the body, irregular heartbeat, problems with breathing. With prolonged use of flax seeds in quantities exceeding the daily norm, the nervous system is affected. In the most severe cases, cyanide poisoning can be fatal.

The EFSA report compared cyanide content in various products. Flax seeds, as well as bitter almonds, are called one of the "foods with the highest cyanide levels." The amygdalin they contain decomposes to release hydrogen cyanide, an aqueous solution of which is known as hydrocyanic acid.

But not everything is as bad as it may seem at first glance. Scientists only warn about the sad consequences in the case of abuse of flax seeds and talk about the worst-case scenario. In fact, you can include this product in your diet, but in reasonable doses. Mayo Clinic.org writes that no cyanide-related side effects were reported after taking flaxseed supplements at the recommended daily highs.

The Food and Drug Administration (FDA) reports that most manufacturing processes wash out this type of cyanide because cyanogenic glycosides break down when exposed to elevated temperatures. Culinary processing (cooking in bread, muffins, etc.) also helps to eliminate the risk of poisoning.

To the question of whether flax seeds lose their useful properties during heat treatment, there is a clear answer. In studies conducted in the United States, the following conditions were set: flaxseeds were crushed and added to food using an oven at a temperature of at least 150 °C. The shortest baking time was 15 minutes, and the longest was 3 hours.

In all cases, linolenic acid (Omega-3) contained in linseed oil remained stable and unchanged. Currently, flax seeds are considered a functional food product, the functional ingredients of which have a proven physiological effect on the human body.

Conclusions:

1. The composition of flaxseed determines its value as a dietary product, but you should not eat seeds directly from the plant that have not been properly processed.
2. It is recommended that an adult consumes no more than 10 grams of flaxseed or flaxseed flour daily to avoid possible side effects.

List of literature:

1. Duke J. A. Handbook of Phytochemical Constituents of GRAS Herbs and other Economic Plants. Boca Raton: CRC Press, 1992.
2. Bhatta R. S. Flaxseed in Human Nutrition. Ed. by S. C. Cunnane and L. U. Thompson. AOAC Press. Champaign, IL., 1995, p. 22-42.
3. Kaprelyants L.V., Shvets N. A. Biochemical characteristics of flax seed lipids. H.A. Zernovye produkty i kombikorma [Grain products and mixed feed].- 2002.- №1.
4. Sarkisova V., Xegay R., Numonova A. ENDOCRINE CONTROL OF THE DIGESTION PROCESS. GASTROINTESTINAL ENDOCRINE CELLS //Science and innovation. – 2022. – T. 1. – №. D8. – C. 582-586.
5. Sarkisova B., Abdurakhmanova K. The role of hormonal drugs in the treatment of endometrial hyperplastic processes and in particular in uterine fibroids. 2014. Vol. 1, no. 01, pp.167-168..
6. Mamatkulovna V. A., Vladimirovna M. E., Mammadov A. N. Properties of Strains of Staphylococcus Aureus Taken From People in Rural Areas in Winter Conditions

//INTERNATIONAL JOURNAL OF HEALTH SYSTEMS AND MEDICAL SCIENCES. – 2022. – T. 1. – №. 5. – С. 92-94.

7. Sarkisova V V, Ismatova M Sh, Rakhmatova F U, Xegay R Adenomyosis as an Independent Unit of Dysfunction of the Endometrium and Uterine Myometrium
8. Ismatova M Sh, Rakhmatova F U, Rakhmatova S U Factors Influencing the Threshold of Pain and Tactile Sensitivity
9. in Medical Students
10. Charos Kuchimova, Marguba Ismatova, Farangiz Yuldasheva, Tolib Turaev. FEATURES OF DRUG ADDICTION AND PHASE DEPRESSIVE DISORDERS IN DYSTHYMIC DISORDERS. JOURNAL OF BIOMEDICINE AND PRACTICE VOLUME 6, ISSUE 1, Tashkent 2021, p. 270.
11. Karabayev Aminjon Gadaevich, Karabayeva Marjona Aminjonovna, Xudoyarova Dildora Raximovna. Study of vegetative reactivity of pregnant women with normoblastic normochrom ic hematopoiesis. /Polish science journal. -2021.-№8.-С.36-55.
12. Ismatova M Sh, Rakhmatova F U WAYS OF ORGANIZING INTERACTIVE LESSONS IN PHYSIOLOGY IN MEDICAL UNIVERSITIES JournalNX- A Multidisciplinary Peer Reviewed Journal ISSN No: 2581 - 4230 VOLUME 9, ISSUE 9, Sep. -2023.
13. Rakhmatova F U, Narzullayeva U.R PATHOGENETIC MECHANISMS OF DISORDERS IN THE HEMOSTASIS SYSTEM OBSERVED IN PATIENTS INFECTED WITH COVID-19 Журнал IEJRD - International Multidisciplinary Journal, vol. 7, no. ICMEI, p. 3, 13 Feb. 2023.