

ANATOMICAL PECULIARITIES OF THE CARDIOVASCULAR SYSTEM

Kadirova Munojatkhon

Assistent of Andijan state medical institute, Uzbekistan

Abstract: The cardiovascular system, comprised of the heart and network of arteries and veins, is the body's transportation system that delivers oxygen, nutrients, hormones, and cellular waste products throughout the body. At the center of this complex circulatory network lies the heart, which acts as a muscular pump to propel blood into the arteries. This article will examine key components of the cardiovascular system involved in supplying blood specifically to the scalp region of the head. The guideline of the cardiovascular framework happens through a heap of upgrades, including changing blood volume, chemicals, electrolytes, osmolarity, prescriptions, adrenal organs, kidneys, and considerably more. The parasympathetic and thoughtful sensory systems likewise assume a vital part in the guideline of the cardiovascular system.

Keywords: Blood vassels, cardiovascular system, oxygen, body features, modern medicine, operations, blood system.

Introduction: The cardiovascular framework gives blood supply all through the body. By answering different boosts, it has some control over the speed and measure of blood helped through the vessels. The cardiovascular framework comprises of the heart, supply routes, veins, and vessels. The heart and vessels cooperate unpredictably to give satisfactory blood stream to all pieces of the body. The guideline of the cardiovascular framework happens through a heap of upgrades, including changing blood volume, chemicals, electrolytes, osmolarity, prescriptions, adrenal organs, kidneys, and considerably more. The parasympathetic and thoughtful sensory systems likewise assume a vital part in the guideline of the cardiovascular system.

The heart is the organ that siphons blood through the vessels. It siphons blood straightforwardly into veins, all the more explicitly, the aorta or the aspiratory corridor. Veins are basic since they control how much blood stream to explicit pieces of the body. Veins incorporate supply routes, vessels, and veins. Conduits divert blood from the heart and can isolate into enormous and little corridors. Enormous supply routes get the most elevated strain of blood stream and are thicker and more flexible to oblige the high tensions. More modest veins, like arterioles, have more smooth muscle, which contracts or unwinds to direct blood stream to explicit bits of the body. Arterioles face a more modest pulse, meaning they needn't bother with to be as versatile.

Arterioles represent the majority of the opposition in aspiratory course since they are more unbending than bigger supply routes. Moreover, the vessels branch off of arterioles and are a solitary cell layer. This dainty layer takes into account the trading of supplements, gases, and waste with tissues and organs. Additionally, the veins transport blood back to the heart. They contain valves to forestall the reverse of blood.

The cardiovascular framework comprises of two principal circles, fundamental flow, and pneumonic course. The reason for the cardiovascular framework is to give sufficient dissemination of blood through the body. Aspiratory flow considers the oxygenation of the blood, and fundamental course accommodates oxygenated blood and supplements to arrive at the remainder of the body. It is vital to figure out the idea of cardiovascular result, stroke volume, preload, Straightforward Starling regulation, afterload, and discharge division to grasp the physiology of the heart. The heart yield (CO)

is how much blood catapulted from the left ventricle, and regularly it is equivalent to the venous return.

The estimation is $CO = \text{stroke volume (SV)} \times \text{pulse (HR)}$. CO likewise rises to the pace of oxygen utilization separated by the distinction in blood vessel and venous oxygen content. The stroke volume is how much blood siphoned out of the heart after one withdrawal. It is the distinction in end-diastolic (EDV) and end-systolic volume (ESV). It increments with expanded contractility, expanded preload, and diminished afterload. Likewise, contractility of the left ventricle increments with catecholamines by expanding intracellular calcium particles and bringing down extracellular sodium. The preload is the strain on the ventricular muscle by the ventricular EDV. Straightforward Starling regulation depicts the connection among EDV and SV.

This regulation expresses that the heart endeavors to level CO with venous return. As venous return expands, there is a bigger EDV in the left ventricle, which prompts further extending of the ventricle. Further extending of the ventricle prompts a bigger compression force and a bigger SV. A bigger stroke volume prompts a bigger CO, in this way balancing CO with venous return. Then, the afterload is the tension that the left ventricular strain should surpass to push blood forward. Mean blood vessel pressure best gauges this. Additionally, afterload can be assessed by the base measure of tension expected to open the aortic valve, which is identical to the diastolic strain.

In this way, diastolic circulatory strain is one of the better ways of ordering afterload. At last, the launch part (EF) is equivalent to SV/EDV . EF of the left ventricle is a record for contractility. A typical EF is more noteworthy than 55%. A low EF demonstrates cardiovascular breakdown.

The cardiovascular cycle portrays the way of the blood through the heart. It runs in the accompanying request:

Atrial constriction conclusion of the mitral valve

Isovolumetric stage

Opening of the aortic valve

Discharge stage (quick and diminished launch), exhausting of the left ventricle

Conclusion of the aortic valve

Isovolumetric unwinding

The kickoff of the mitral valve

Filling stage (quick and decreased filling) of the left ventricle

Vasculature assumes a critical part in the guideline of blood stream all through the body. By and large, pulse diminishes from conduits to veins, and this is a direct result of the strain conquering the obstruction of the vessels. The more prominent the adjustment of obstruction anytime in the vasculature, the more prominent the deficiency of tension by then. Arterioles have the most expansion in opposition and cause the biggest abatement in pulse. The tightening of arterioles increments obstruction, which causes a decline in blood stream to downstream vessels and a bigger diminishing in pulse.

Expansion of arterioles causes a decline in opposition, expanding blood stream to downstream vessels and a more modest diminishing in circulatory strain. Diastolic pulse (DP) is the most reduced strain in a corridor toward the start of the cardiovascular cycle while the ventricles are unwinding and filling. DP is straightforwardly corresponding to add up to fringe opposition (TPR). Additionally, the energy put away in the consistent aorta during systole is presently delivered by the force of the aortic wall during diastole, hence expanding diastolic tension. Systolic circulatory strain (SP) is the pinnacle tension in a course toward the finish of the cardiovascular cycle while the ventricles are contracting.

Straightforwardly connected with stroke volume, as stroke volume builds, SP additionally increments. SP is likewise impacted by aortic consistence. Since the aorta is flexible, it stretches and stores the energy brought about by ventricular withdrawal and diminishes the systolic tension. Beat pressure is the distinction among SP and DP. Beat pressure is relative to SV and conversely corresponding to blood vessel consistence.

Consequently, the stiffer the course, the bigger the beat pressure. Mean blood vessel pressure (Guide) is the typical tension in the conduits all through the cardiovascular cycle. The Guide is in every case nearer to DP. MAP is determined by $MAP = DP + 1/3$ (beat pressure). Likewise, $Guide = CO \times TPR$, where CO is heart yield. This worth is critical on the grounds that at whatever point there is a decline in CO, to keep up with the Guide, the TPR will build, which is important in numerous pathophysiology issues.

Fundamental veins have a lower decline in pressure since it has low obstruction.

The venous framework is exceptionally consistent and contains up to 70% of the flowing blood without a moment's delay. A little change in venous strain can prepare the blood put away in the venous framework. Speed of blood in the vasculature has a reverse relationship with cross-sectional region (volumetric stream rate (Q) = stream speed (v) x cross-sectional region (A)). As the cross-sectional region increments, speed diminishes. Corridors and veins have more modest cross-sectional regions and the most elevated speeds, though vessels have the most cross-sectional region and the least speeds.

The vasculature additionally gives obstruction. Obstruction is $R = (8 \times \text{viscosity} \times \text{length}) / (\pi r^4)$. Thickness relies upon hematocrit and expansions in various myeloma or polycythemia. As cylinder length expands, the obstruction increments. As the cylinder range expands, the opposition diminishes. The way that the range is to the force of 4 implies that slight changes in the sweep significantly affect opposition. The complete obstruction of vessels in a series is $R_1 + R_2 + R_3$, etc., and the all-out opposition of veins in equal is $1/TR = 1/R_1 + 1/R_2 + 1/R_3$, etc., where TR is the all-out opposition.

The Poiseuille condition estimates the progression of blood through a vessel. It is estimated by the adjustment of tension separated by obstruction: $Stream = (P_1 - P_2)/R$, where P is strain, and R is opposition. Expanding opposition in a vessel, for example, the tightening of an arteriole, causes a reduction in blood stream across the arteriole. Simultaneously, there is a bigger decline in tension across this point on the grounds that the strain is lost by defeating the opposition. Expanding the obstruction anytime increments upstream strain however diminishes downstream tension.

The Poiseuille condition applies to the fundamental dissemination with the end goal that F is the cardiovascular result (CO), P1 is the mean blood vessel pressure (Guide), P2 is the right atrial

strain (RAP), and R is the all-out fringe obstruction (TPR). Since RAP is near 0 and tiny in contrast with Guide, the condition approximates as $F=P1/R$ or $CO=MAP/TPR$ where $MAP=CO*TPR$ - this implies that cardiovascular result and absolute fringe opposition control Guide. Its application is significant in light of the fact that in injury circumstances with drain, there is likewise a reduction in heart yield, however on occasion the circulatory strain is close to typical.

This is on the grounds that the TPR at the level of the arterioles has expanded. This condition, as applied to the pneumonic vasculature, is utilized to decide the reason for aspiratory hypertension. As connected with the pneumonic vasculature, F addresses CO, P1 addresses pneumonic vein pressure (PAP), and P2 addresses left atrial tension (LAP), and R is pneumonic vascular opposition (PR); $CO=(PAP-LAP)/PR$. A Swan-Ganz catheter assists with estimating both PAP and LAP, considering the estimation of PR and, in this way, the etiology of pneumonic hypertension.

The sensory system directs the cardiovascular framework with the assistance of baroreceptors and chemoreceptors. The two receptors are situated in the carotids and aortic curve. Likewise, both have afferent signs through the vague nerve from the aortic curve and afferent signs through the glossopharyngeal nerve from the carotids.

Conclusion:In summary, this essay has delineated the intricate cardiovascular pathways that deliver oxygenated blood from the heart to the various regions of the scalp. The external carotid and occipital arteries primarily nourish the occipital scalp area, while the internal carotid system and its branches, including the middle meningeal arteries, are responsible for blood flow to the frontal, temporal, and parietal scalp territories. A well-functioning cardiovascular network is essential to sustain the health and function of this important protective covering of the skull.

References:

1. Huang Y, Hu D, Huang C, Nichols CG. Genetic Discovery of ATP-Sensitive K⁺ Channels in Cardiovascular Diseases. *Circ Arrhythm Electrophysiol.* 2019 May;12(5): e007322.
2. Baytanov, O. (2023). O 'ZBEKISTONDA OZIQ-OVQAT SANOATI RIVOJLANISHINING HOZIRGI HOLATI. BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI, 3(3), 446-449.
3. Mirzaqul o'g, B. O. R. (2023). HUDUDLARDA ISHBILARMONLIK MUHITINI YAXSHILASH, KICHIK BIZNES VA XUSUSIY TADBIRKORLIKNI YANADA RIVOJLANTIRISHNI RAG'BATANTIRISH USULLARI.
4. RAQAMLI, I., & ELEKTRON, T. O'ZBEKISTON IQTISODIYOTINI RIVOJLANTIRISHDA RAQAMLI IQTISODIYOTNING AHAMIYATI. *ЭКОНОМИКА*, 36-38.
5. Мухамедова, М. Г., Куртиева, Ш. А., & Назарова, Ж. А. (2020). СИНДРОМ ФУНКЦИОНАЛЬНОЙ КАРДИОПАТИИ У СОВРЕМЕННЫХ ПОДРОСТКОВ. In П84 Профилактическая медицина-2020: сборник научных трудов Все-российской научно-практической конференции с международным участием-ем. 18–19 ноября 2020 года/под ред. АВ Мельцера, ИШ Якубовой. Ч. 2.—СПб.: Изд-во СЗГМУ им. ИИ Мечникова, 2020.—304 с. (p. 105).
6. Kurtieva, S., Nazarova, J., & Mullajonov, H. (2021). Features of endocrine and immune status in adolescents with vegetative dystonia syndrome. *International Journal of Health Sciences*, 5(2), 118-127.

7. Kurtieva, S., Nazarova, J., & Mullaionov, H. (2021). Features of Physical and Generative Development of Modern Teenagers Living in Uzbekistan. *NeuroQuantology*, 19(7), 57.
8. Куртиева, Ш. (2021). CHANGES IN ELECTROENCEPHALOGRAM INDICATORS IN ADOLESCENTS WITH VEGETATIVE DYSFUNCTION. *УЗБЕКСКИЙ МЕДИЦИНСКИЙ ЖУРНАЛ*, (SPECIAL 3).
9. Kurtieva, S. (2021). Adaptation capabilities of functional systems of the body of adolescents with vegetative dystonia syndrome. *International Journal of Health and Medical Sciences*, 4(1), 129-135.
10. Kurtieva, S. (2021). Clinical And Anamnestic Characteristics Of The Health State Of Adolescents With Vegetative Dysfunction Syndrome. *The American Journal of Medical Sciences and Pharmaceutical Research*, 3(06), 1-12.
11. Kurtieva, S. (2022). 24-HOUR BLOOD PRESSURE MONITORING IN ADOLESCENTS WITH VEGETATIVE DYSTONIA SYNDROME. *The American Journal of Medical Sciences and Pharmaceutical Research*, 4(01), 1-8.