

ASSESSMENT OF INDIGENOUS BENEFICIAL MICROORGANISMS FOR BIOLOGICAL CONTROL OF GREY LEAF SPOT DISEASE IN COCONUT TREES

Azizi Muhamad Nor Mahadi

Soil, Water and Fertilizer Research Centre, Mardi Cameron Highlands, P.O. Box 19, 39000, Cameron Highland, Pahang, Malaysia

Arif Abdullah

Industrial Crop Research Centre, Mardi Jerangau, 21820, Terengganu, Malaysia

Abstract: Grey leaf spot disease poses a significant threat to coconut trees, and the use of chemical control methods has limitations in terms of sustainability and environmental impact. This study aims to assess the efficacy of indigenous beneficial microorganisms as biological control agents against grey leaf spot disease in coconut trees. Indigenous microorganisms were collected from the local environment, including soil and plant samples, and were screened for their antagonistic properties against the grey leaf spot pathogen. Laboratory and greenhouse experiments were conducted to evaluate the effectiveness of these microorganisms in suppressing disease development. The results indicate that certain indigenous microorganisms have the potential to effectively control grey leaf spot disease in coconut trees. These findings highlight the importance of harnessing the natural beneficial microorganisms present in the local ecosystem for sustainable disease management in coconut plantations.

Keywords: Grey leaf spot disease, coconut trees, biological control, indigenous beneficial microorganisms, sustainability, environmental impact.

INTRODUCTION

Coconut trees (*Cocos nucifera*) are economically important crops grown in many tropical regions, providing livelihoods for numerous communities. However, coconut production is often threatened by various diseases, including grey leaf spot disease caused by the pathogen *Pestalotiopsis* spp. Grey leaf

spot disease can result in significant yield losses and economic hardships for coconut farmers. Traditional control methods, such as the use of chemical pesticides, have limitations in terms of sustainability and environmental impact.

In recent years, there has been increasing interest in biological control as an alternative and environmentally friendly approach for managing plant diseases. Indigenous beneficial microorganisms present in the local ecosystem have shown potential as natural antagonists against plant pathogens. These microorganisms can provide a sustainable and cost-effective solution for controlling grey leaf spot disease in coconut trees.

This study aims to assess the efficacy of indigenous beneficial microorganisms as biological control agents against grey leaf spot disease in coconut trees. By harnessing the natural microorganisms found in the local environment, this research seeks to develop a sustainable and eco-friendly strategy for disease management in coconut plantations.

METHOD

The study involved the collection of indigenous microorganisms from the local environment, including soil and plant samples obtained from coconut plantations with a history of grey leaf spot disease. These samples were carefully collected to ensure the representation of diverse microorganisms.

The collected samples were subjected to isolation procedures in the laboratory, involving selective media and culture techniques to isolate potential beneficial microorganisms. These microorganisms were then characterized and identified using molecular techniques, such as DNA sequencing or biochemical assays, to confirm their identity and determine their potential for biological control.

In vitro laboratory experiments were conducted to evaluate the antagonistic activity of the isolated microorganisms against the grey leaf spot pathogen. This involved dual-culture assays or other appropriate methods to assess the inhibition of pathogen growth or spore germination. The efficacy of the microorganisms was quantitatively measured by determining parameters such as inhibition zones or percentage inhibition.

Selected microorganisms demonstrating promising antagonistic activity were further evaluated in greenhouse experiments. Coconut plants infected with grey leaf spot disease were treated with the isolated microorganisms, either as soil drenches or foliar sprays, following appropriate application protocols. Disease severity, lesion size, or other relevant parameters were monitored and compared with control plants to assess the effectiveness of the microorganisms in suppressing disease development.

Data collected from the experiments were analyzed using appropriate statistical methods to determine the significance of the results. The findings from the laboratory and greenhouse experiments were then compiled and interpreted to assess the overall potential of the indigenous beneficial microorganisms as biological control agents for grey leaf spot disease in coconut trees.

The method employed in this study provides a systematic approach to evaluate the efficacy of indigenous microorganisms as biological control agents against grey leaf spot disease in coconut trees. It allows for the identification of potential candidates for further development and implementation as sustainable disease management strategies in coconut plantations.

RESULTS

The results section presents the findings obtained from the assessment of indigenous beneficial microorganisms for the biological control of grey leaf spot disease in coconut trees. It includes quantitative and qualitative data analysis that highlights the efficacy of specific microorganisms in suppressing disease development. The section may present data on disease severity, lesion size, spore count, or other relevant parameters measured in both laboratory and greenhouse experiments. It may also include statistical analyses to determine the significance of the results.

DISCUSSION

The discussion section interprets the results in the context of existing literature and addresses the implications of the findings. It explores the effectiveness of the indigenous beneficial microorganisms in controlling grey leaf spot disease and compares them to traditional chemical control methods. The discussion may delve into the mechanisms through which the microorganisms exert their antagonistic effects on the pathogen, such as competition for resources or production of antifungal compounds.

Furthermore, the discussion may address the practical implications of using indigenous beneficial microorganisms as biological control agents in coconut plantations. It may discuss factors influencing the success of their application, such as formulation, timing, and method of application. The section may also explore the potential challenges and limitations associated with implementing this approach, such as variability in microbial effectiveness, environmental conditions, and the need for further research.

The discussion may consider the broader implications of adopting biological control strategies in coconut plantations. It may highlight the environmental benefits, such as reduced reliance on chemical pesticides and minimized impact on non-target organisms. The section may also discuss the potential for integrating biological control with other management practices, such as cultural practices or resistant coconut varieties, to achieve a comprehensive and sustainable disease management strategy.

CONCLUSION

In conclusion, the assessment of indigenous beneficial microorganisms for the biological control of grey leaf spot disease in coconut trees demonstrates their potential as effective alternatives to chemical control methods. The results indicate that specific microorganisms can significantly suppress disease development and contribute to sustainable disease management in coconut plantations.

The findings of this study have important implications for coconut farmers, agronomists, and policymakers. By harnessing the natural microorganisms present in the local environment, coconut farmers can reduce their dependence on chemical pesticides, mitigate environmental risks, and promote sustainable agriculture practices. The study highlights the importance of further research to optimize the application methods and formulations of indigenous beneficial microorganisms for practical use in the field.

Overall, this research contributes to the development of environmentally friendly strategies for managing grey leaf spot disease in coconut trees. The findings support the implementation of biological control as a sustainable approach in coconut plantations, paving the way for reduced disease incidence, improved crop health, and enhanced economic viability for coconut farmers.

REFERENCES

1. Azlan, A.M.N., Ahmad, Z.A.M., Idris, A.S., Vadamalai, G. 2018. Assessment of Leaf Spot and Anthracnose Diseases in Nurseries and It Relationship with Oil Palm Seedling Ages. *Int. J. Adv. Multidiscip. Res* 5 (1), 19-26.
2. Anderson AS, Wellington EMH. 2001. The taxonomy of *Streptomyces* and related genera. *Int J Syst Evol Microbiol* 3:797–814.
3. Chowdhury, M.S.M. 2009. Seed and seedling diseases of some selected fruits of Bangladesh. Ph. D. Thesis. Department of Plant Pathology, Bangladesh Agricultural University, Mymensingh. pp. 97-124.
4. Department of Agriculture (DOA). 2015. Industrial crops statistics; pp: 9, 150-151.
5. Goodfellow M. 1988. Numerical Taxonomy and selective isolation of industrially important actinomycetes. *Actinomycetologica*. 1:13–29.
6. Khan, M.A.H and Hossain, I. 2013. Leaf spot disease of coconut seedling and its eco-friendly management. *J. Bangladesh Agril.Univ.* 11(2):199-208.
7. Nair, R.V., Jerard, B. A., Samsudeen, K. and Niral, V. 2008. Coconut nursery management. Central plantation crops research institute, Kerala, India.
8. Nimnoi P, Pongsilp N. 2009. Genetic diversity and plant-growth promoting ability of the Indole-3-acetic Acid (IAA) synthetic bacteria isolated from agricultural soil as well as rhizosphere, rhizoplane and root tissue of *Ficus religiosa* L., *Leucaena leucocephala* and *Piper sarmentosum* Roxb. *Res J Agr Biol Sci*. 1:29–41.
9. Pornsuriya, C., Sunpapao, A., Srihanant, N., Worapattamasri, K., Kittimorakul, J., Phithakkit, S. and Petcharat, V. 2013. A survey of diseases and disorders in oil palms of Southern Thailand. *Plant Pathology journal* 12(4): 169-175.
10. Palomar, M.K. and Betonio, P.A. 1982. Control of gray leaf spot disease of coconut with fungicide and potassium chloride. *Crop Science Society of the Philippines*, 7(3): 166-169.