

*Abdusattarov A.A.**Preparation of Family Doctors**Andijan State Medical Institute, Republic of Uzbekistan***TREATMENT AND DIAGNOSIS OF FUNCTIONAL DYSPEPSIA IN PEOPLE WITH DIABETES 2 TYPES****Keywords:** dyspepsia syndrome, type 2 diabetes mellitus, *Helicobacter pylori*.

Diabetes mellitus (DM) is a group of metabolic diseases characterized by high rates of morbidity [1, 2] and systemic complications involving various organs and systems, including the gastrointestinal tract [1-3]. Symptoms of dyspepsia are among the most common complaints related to digestive disorders and occur in 25% of the population [4, 5]. The concept of dyspepsia has been repeatedly revised and clarified. According to the Rome Criteria II (1999), dyspepsia is understood as a feeling of pain or discomfort, early satiety, overflow localized in the epigastric region, as well as bloating or nausea [6]. Based on the assessment of the leading clinical manifestation, dyskinetic and ulcer-like variants of dyspepsia syndrome are distinguished [6-8].

In the dyskinetic variant, symptoms of discomfort, early satiety, feelings of fullness, swelling in the epigastric region and nausea predominate, while in the ulcer-like case, epigastric pain dominates. A number of studies have shown that symptoms of dyspepsia are observed in patients with type 2 diabetes more often than in patients without impaired carbohydrate metabolism [9]. The results of studies of dyspepsia syndrome in patients depending on the compensation of carbohydrate metabolism [9, 10], as well as infection with *Helicobacter pylori* are contradictory [11, 12].

The purpose of the study. To study the prevalence, clinical manifestations of dyspepsia syndrome and factors associated with its occurrence in patients with type 2 diabetes.

Materials and methods. A single-center, single-stage observational descriptive comparative study was conducted. The stages of the work were carried out with the voluntary consent of the patients. After a thorough examination, taking into account the inclusion and exclusion criteria, A group of patients has been formed for further research. Inclusion criteria: Type 2 diabetes. Exclusion criteria: conditions and diseases in the stages requiring urgent intervention, impaired cognitive function, the use of drugs from the group of nonsteroidal anti-inflammatory drugs, antibiotics, theophylline, cardiac glycosides, potassium and iron preparations that can provoke symptoms of dyspepsia during the last 12 months. The formed group consisted of 107 patients with type 2 diabetes (38.3% of women, 61.7% of men) aged 54.2 ± 10.1 years. The duration of type 2 diabetes ranged from 0 to 20 years (on average, 6.9 ± 5.81 years).

Compensation of carbohydrate metabolism by the level of glycated hemoglobin (HbA1c) averaged $7.3 \pm 1.21\%$. Treatment of hyperglycemia included oral hypoglycemic drugs, insulin therapy, or combination treatment. None of the patients included in the study associated the presence of dyspepsia syndrome with the prescribed treatment. There were no differences in the frequency of medication intake in the groups, depending on the presence or absence of dyspepsia. The diagnosis of type 2 diabetes was confirmed by assessing the glycemic profile and HbA1c level [1, 2]. The WHO classification (1999) was used to formulate the diagnosis of type 2 diabetes.

All patients were clinically examined in detail with an assessment of complaints, anamnesis, objective examination of laboratory parameters of carbohydrate metabolism, clinical and biochemical screening of diabetes complications. All patients underwent endoscopic examination of the upper gastrointestinal mucosa with histological detection of *Helicobacter pylori* in the gastric mucosa. A

detailed examination revealed dyspepsia syndrome according to the Roman Criteria II [6] in 76 patients. Of these, 32 patients (average age 56 ± 8.65 years, 27.3% of women, 72.7% of men) had organic gastrointestinal diseases that could cause this syndrome.

In 44 patients (57.7 ± 9.81 years old, 45.5% of women, 54.5% of men), dyspepsia syndrome could not be explained by an organic pathology of the gastrointestinal tract. 31 patients with type 2 diabetes (49.3 ± 8.22 years old, 54.8% of men, 45.2% of women) had no complaints of dyspepsia symptoms. A group of patients with functional dyspepsia was also formed by continuous sampling for comparative studies ($n=33$; 41.2 ± 10.61 years, 16.1% of women, 83.9% of men). Statistical data processing was carried out using SPSS 11.5 software packages.

For quantitative data, the mean (M) and standard deviation (SD) were calculated, the results are presented as $M\pm SD$. The Mann-Whitney U—test was used for paired intergroup comparison of indicators, and the Pearson correlation coefficient was used to determine the relationship between the studied features. The odds ratio (OR) with 95% confidence intervals (95% CI) was calculated using contingency tables. The critical significance level when testing statistical hypotheses (p) was assumed to be less than 0.05.

Results and discussion. Dyspepsia syndrome was detected in 76 (71%) of 107 patients with type 2 diabetes included in the study. According to other studies, the incidence of dyspepsia symptoms in patients with type 2 diabetes was higher than in the general population [9, 10, 13], and reached 18.2-68%, depending on the criteria of dyspepsia syndrome and the clinical characteristics of the examined patients with type 2 diabetes. In contrast to patients with type 2 diabetes without dyspepsia syndrome, patients with type 2 diabetes with dyspepsia syndrome had a 2-fold longer duration of diabetes due to individuals who were observed for 5 years or more (8.7 ± 6.3 and 4.3 ± 3.7 years, respectively; $p=0.001$).

Patients with dyspepsia had higher HbA1c levels (7.5 ± 1.3 and $7\pm 0.79\%$; $p=0.03$) and lower glomerular filtration rate (GFR) (100.3 ± 24.83 and 113.2 ± 15.71 ml/min; $p=0.013$). In the type 2 diabetes group with dyspepsia, diabetic retinopathy was 2.6 times more common, diabetic neuropathy was 2 times more common, coronary heart disease was 3.9 times more common ($p<0.006$), H. pylori infection was 2.1 times higher (58 and 27%; $p=0.007$). Thus, dyspepsia syndrome was more often detected in patients with type 2 diabetes with a longer duration of the disease, poorer glycemic parameters, against which complications of diabetes (retinopathy, neuropathy, coronary heart disease) are formed, in patients with H. pylori infection.

Of the 76 patients with type 2 diabetes with dyspepsia, 42.1% had organic gastrointestinal diseases that can explain the presence of dyspepsia syndrome: 34.4% had biliary tract pathology, 25% had gastric or duodenal ulcer, 21.9% had gastroesophageal reflux disease, 18.7% had gastric and/or duodenal erosion guts. In the remaining 57.9%, dyspepsia could not be explained by concomitant gastrointestinal diseases. As can be seen in the figure, in patients with type 2 diabetes with dyspepsia syndrome, which cannot be explained by the organic pathology of the gastrointestinal tract, the dyskinetic variant prevailed, which corresponds to the data of other researchers. [10, 13].

A feeling of fullness was observed in 70% of the surveyed, and discomfort in the epigastrium was observed in 61%. In patients with type 2 diabetes with dyspepsia and organic gastrointestinal pathology, which causes the symptoms of dyspepsia, as in patients with functional dyspepsia, the ulcer-like variant of dyspepsia with a predominance of epigastric pain prevailed, which was observed in 59 and 73% of cases, respectively. The frequency of detection of early saturation and overflow in patients with type 2 diabetes with dyspepsia, which cannot be explained by organic diseases of the gastrointestinal tract, was significantly higher ($p<0.004$) than in other groups. Patients with type 2 diabetes with dyspepsia syndrome caused by organic gastrointestinal pathology, compared with patients with dyspepsia, which cannot be explained by concomitant gastrointestinal diseases, had

similar clinical (duration of diabetes, body mass index - BMI, complications of diabetes), anthropometric indicators (gender and age), H. pylori infection rate and laboratory parameters.

Attention was drawn to the presence of decompensated carbohydrate metabolism (HbA1c levels of 7.07 ± 1.09 and $7.5 \pm 1.3\%$; $p=0.019$) and hypercholesterolemia (5.6 ± 0.96 and 5.8 ± 1.00 mmol/l; $p=0.72$) in both groups. Compared with patients with functional dyspepsia, patients with type 2 diabetes with dyspepsia were older (57.0 ± 9.84 and 41.2 ± 10.61 years; $p=0.001$), had a higher BMI (29.8 ± 3.30 and 25.6 ± 1.60 kg/m²; $p=0.001$), they were more likely to have coronary heart disease (75 and 18.0%; $p=0.005$), lipid metabolism disorders were more common (total cholesterol levels 5.8 ± 1.00 and 5.4 ± 1.04 mmol/l; $p=0.04$), renal GFR was lower (100.3 ± 24.82 and 118.5 ± 15.71 ml/min; $p=0.0001$). A comparison of the selected groups allowed us to conclude that dyspepsia syndrome has different manifestations depending on the clinical characteristics (patients with type 2 diabetes with or without concomitant gastrointestinal diseases, patients with functional dyspepsia).

To identify the factors associated with dyspepsia syndrome, clinical features (BMI, duration of the disease, the presence of complications of diabetes - diabetic retinopathy, diabetic neuropathy, nephropathy, non—alcoholic fatty liver disease, coronary heart disease), anthropometric parameters (gender and age), laboratory parameters were analyzed in patients with type 2 diabetes without organic gastrointestinal pathology. (basal blood glucose, fasting blood glucose, HbA1c concentration, total cholesterol, alanine and aspartic aminotransferases, creatinine, GFR), H. pylori infection.

According to the data presented, associations of dyspepsia syndrome with the duration of type 2 diabetes, the presence of complications such as diabetic retinopathy and neuropathy, and coronary heart disease have been identified. Consequently, dyspepsia syndrome is formed along with other complications of diabetes.

The above makes it possible to consider dyspepsia syndrome as one of the complications of type 2 diabetes. Dyspepsia syndrome was also associated with the presence of H. pylori infection and the age of patients. Decompensation of carbohydrate metabolism in patients with type 2 diabetes contributes to the formation of various complications of diabetes [1, 2], including dyspepsia syndrome. In addition to dyspepsia syndrome as a whole, some of its symptoms were also associated with some characteristics of the underlying disease. In the table. 2 showed that the symptom of overflow in the epigastrium was associated with the presence of diabetic retinopathy and neuropathy, coronary heart disease, HbA1c levels, and fasting blood glucose levels.

The symptom of discomfort was associated with the duration of type 2 diabetes, the presence of diabetic neuro- and retinopathy, coronary heart disease, H. pylori infection, fasting blood glucose, basal glycemia, and HbA1c. The nausea symptom was associated with the presence of diabetic retinopathy. These results allow us to consider dyspepsia syndrome in patients with type 2 diabetes as a consequence of hyperglycemic syndrome and confirm the view that dyspepsia syndrome should be considered as one of the gastroenterological complications of type 2 diabetes, and the disease itself as a multiple organ pathology.

Conclusion. Dyspepsia syndrome in type 2 diabetes is observed in 71% of cases; in 42.3% it can be explained by organic diseases of the gastrointestinal tract, and in 57.7% there is no connection with the pathology of the digestive system. Dyspepsia syndrome in patients with type 2 diabetes, which cannot be explained by organic diseases of the gastrointestinal tract, is manifested mainly by the dyskinetic variant, while in patients with type 2 diabetes with organic diseases of the gastrointestinal tract and in patients with functional dyspepsia, the ulcer-like variant prevails. Dyspepsia syndrome in patients with type 2 diabetes, which cannot be explained by gastrointestinal diseases, is associated with the duration of carbohydrate metabolism disorders, the presence of diabetic retinopathy and neuropathy, coronary artery disease, H. pylori and the age of patients.

Some manifestations of dyspepsia syndrome (a symptom of overflow and discomfort in the epigastrium), which cannot be explained in patients with type 2 diabetes by organic gastrointestinal diseases, are also associated with indicators of carbohydrate metabolism: a symptom of overflow - with the level of HbA1c and fasting blood glucose, a symptom of discomfort — with the level of HbA1c, with basal blood glucose and fasting blood glucose levels.

Literature:

1. Dedov I., Shestakova M.V. Algorithms of specialized medical care for patients with diabetes mellitus. *Diabetes mellitus Supplement to the journal* 2011; 3: 1-72.
2. Dedov I.I., Melnichenko G.A. *Endocrinology: A national guide*. Moscow: GEOTAR-Media 2011; 120-151, 253-329.
3. Leites Yu.G., Galstyan G.R., Marchenko E.V. Gastroenterological complications of diabetes mellitus. *Medical consultation Diligence: Gastroenterology* 2007; 2: 25-32.
4. Harmon R.S., Peura D.A. *Assessment and treatment of dyspepsia*. Gastroenterol Publishing House, 2010; 3 (2): 87-98.
5. Tack J., Talley N.J., Camilleri M. et al. Functional gastroduodenal disorders. *Gastroenterology* 2006; 130 (5): 1466-1479.
6. Talley, N.J., Stangellini, V., Heding, R. S., and others. Functional gastroduodenal disorders. *Gut* 1999; 45 (Supplement II): II37—II42.
7. Keohane J., Quigley E.M. Functional dyspepsia: the role of visceral hypersensitivity in its pathogenesis. *The World of Gastroenterology*, 2006; 12 (17): 2672-2676.
8. Talley N.J., Stangellini V., Heding R.S. and others. Constitutional officials. In: Drosman D.A. (ed.) *Rome II. Functional Gastrointestinal Disorders*, 2nd ed. Allen Press 2000: 299-350.
9. Abid S., Rizvi A., Jahan F. Poor glycemic control is the main factor associated with an increased incidence of gastrointestinal symptoms in patients with diabetes mellitus. *Jay Park, Associate Professor of Medicine*, 2007; 57 (7): 345-349.
10. Oh J.H., Choi M.J., Kang M.I. and others. Prevalence of gastrointestinal symptoms in patients with non-insulin-dependent diabetes mellitus. *Korean International Medical Journal*, 2009; 24 (4): 309-317.
11. Devrajani B.R., S. Shah Z., And Soomro A., Devrajani T. Type 2 diabetes mellitus: a risk factor for *Helicobacter pylori* infection: a case-control clinical trial. *International Diabetes Research*, 2010; 30 (1): 22-26.
12. Gulchelik N. E., Kaya E., Demirbas B. et al. The prevalence of *Helicobacter pylori* in patients with diabetes mellitus and its association with dyspepsia and autonomic neuropathy. *J Endocrinol Invest*, 2005; 28 (3): 214-217.
13. Kuan S., Talley N.J., Jones M.P. and others. Gastrointestinal symptoms and glycemic control in diabetes mellitus: a longitudinal population study. *Eur J Gastroenterol Hepatol*, 2008; 20 (9): 888-897.