

ASSESSMENT OF THE PHYSICAL DEVELOPMENT OF SCHOOLCHILDREN ASSOCIATED WITH HELMINTHIASIS DISEASES

Aida Sh. Kamilova ¹, Djamshid Y. Kamilov ²

1 Assistant of the Department of hygiene of children, adolescents and nutrition of the
Tashkent state Medical university, Tashkent, Uzbekistan.

E-mail:aida.kamilova1988@gmail.com

2 Assistant of the Department of histology and medical biology of the
Tashkent state Medical university, Tashkent, Uzbekistan.

kamilovjamshid3@gmail.com

ABSTRACT

Relevance. Physical development is an indispensable indicator of the health of a growing organism. Subject to general biological laws, it shows a stimulating and depressing effect on environmental factors, including contamination of the living area by various substances that have both. At the same time, the negative effect is not limited only to the moment of exposure, the process of formation of the organism affects not only the future [1,8,9]. The processes of growth and development of the child's body, depending on the environmental conditions that are still little studied, depend on the ecological variability of the habitat and the physics of the growing organism, which can serve as the basis for illuminating the mechanisms of adaptation [2, 3,7].

Age	Standard M±m		Control M±m		Main M±m	
	Boy	Girl	Boy	Girl	Boy	Girl
7	125,14±3,95	124,64±456	125,3±0,78	124.95±0,9 3	123,89±0,4 4	123,00±0,56
8	128,88±4,61	127,89±5,34	128,53±0,59	126,44±0,9 6	127,50±2,5 3	125,50±6,67

Purpose. It consists in a comparative assessment of the main anthropometric indicators of children from 7 to 10 years old in schools in the city of Tashkent.

Research methods and methods. Measurements of anthropometric indicators were carried out in 3,161 children between the ages of 7 and 10, of which 2,651 were educated at secondary school No. 300 in Sergeli district, Tashkent City. The obtained values of anthropometric indicators of physical development of schoolchildren of Sergeli District, Tashkent City were compared with the results of the study of 510 children studying at the secondary school No. 98 in Yunusabad district.

Results. Children from the two districts enter the school with almost the same anthropometric indicators. When we did the research, we divided schoolchildren into two large groups. In this, the controlled group is healthy children, while the main group is the group of children found vomiting. For example, when measured by age, the height indicators were 125.3±0.78 in the 7-year-old boys 'control group, 123.89±0.44 in the main group, and 124.95±0.93 in the 7-year-old girls' control group, and 123.00±0.56 in the main group. 8-year-old boys had 128.53±0.59 in the control group, 127.50±2.53 in the main group, and 126.44±0.96 in the control group of 8-year-old girls, and 126.50±6.67 in the main group (Table 1).

Table 1.

Comparison of the height indicator in the main and control groups of schoolchildren.

9	132,76±5,74	132,04±5,68	132,19±1,16	131,58±1,23	131,55±1,33	130,89±1,99
10	137,42±5,92	137,06±6,32	138,05±0,74	136,89±0,85	136,80±1,46	136,00±1,052

The body weight of shuningdek bolalarning is comparable to the control group of the control groups (Table 2).

Table 2.

The main and control groups for body weighing are trained at the school.

Age	Standard M±m		Control M±m		Main M±m	
	Boy	Girl	Boy	Sex	Boy	Girl
7	25,10±2,96	23,94±2,28	25,08±0,61	23,72±0,78	23,34±0,30	22,18±0,54
8	26,34±2,46	25,14±2,27	27,66±0,85	26,78±0,88	25,75±1,70	24,37±1,05
9	29,82±2,35	27,66±3,47	30,19±1,76	28,83±1,44	28,55±1,12	27,78±1,54
10	32,12±3,04	32,46±3,36	35,31±1,41	33,23±0,96	30,60±0,93	28,75±1,13

The weighing of weights in the trunk, the table top divided into parts, the marks and the analyzing objects must be clear. In 7 cases, the bolalaring control group was 25.08±0.61 ni, the main group was 23.34±0.30 ni; in 7 cases, the bolalaring control group was 23.72±0.78 ni, the main group was 22.18±0.54 ni. Control groups 27.66±0.85 and main groups 25.75±1.70 received 8 points; control groups 26.78±0.88 and main groups 24.37±1.05 received 8 points. 9 of them, the control group was 30.19±1.76, the main group was 28.55±1.12; the control group was 28.83±1.44 and the main group was 27.78±1.54. 10 points above the control group 35.31±1.41 and above the control group 30.60±0.93; above the control group 33.23±0.96 points above the control group 28.75±1.13 points above the control group.

It can be seen from this that the difference in comparison between the control and main groups of children 7 and 8 years old was 1-2 kg, while it turned out that the difference between 9-year-olds is 1 kg, and between the body weight of boys and girls 10 years old is slightly more, that is, 5 kg. Children in the main group were found to have significantly lower body weight than the control group.

Table 3.

Comparison of the index of the circumference of the chest in the main and control groups of schoolchildren

Age	Standard M±m		Control M±m		Main M±m	
	Boy	Girl	Boy	Sex	Boy	Girl
7	62,91±2,27	60,14±2,92	63,94±0,73	62,72±0,58	60,33±0,59	58,55±0,34
8	63,98±2,82	61,94±2,51	64,53±0,67	62,93±1,01	62,25±1,43	58,63±1,07
9	65,90±2,87	62,12±2,33	66,81±1,76	64,04±1,45	64,91±0,69	61,22±1,92
10	69,16±3,03	65,56±3,17	70,13±1,33	66,20±0,87	67,00±0,84	63,25±1,18

The data obtained showed that in children of the main group, negative changes from 3 studied somatometric indicators were mainly due to body weight. The reason for this is characterized by a decrease in appetite and dysfunction of the digestive organs of children suffering from vomiting.

The data obtained showed that in children of the main group, negative changes from 3 studied somatometric indicators were mainly due to body weight. The reason for this is characterized by a decrease in appetite and dysfunction of the digestive organs of children suffering from vomiting.

Conclusion. It can be seen from this that among schoolchildren from 7 to 10 years old, there are significant changes in height, body weight, chest circumference among those in the control and main group. This means that children with vomiting disorders have a downward trend in growth and development compared to healthy children. This does not affect the child's not only height, but also mental and physical development. At the same time, there is also a deficiency of hemoglobin and all vitamins and minerals in the blood, which are involved in the assimilation of Sciences, decreased memory and metabolism.

REFERENCES.

1. Baymamatovich O. B. et al. ANALYSIS OF SOCIO-HYGIENIC INDICATORS IN PATIENTS WITH ORAL CAVITY CANCER. – 2025.
2. Ermatov N. J. et al. Hygienic analysis of the nutritional condition of the employees of textile enterprise. – 2024.
3. Ermatov N. J. I Dr. Eastern awakening: innovation, education, natural and Social Sciences. – 2022. - T. 2. – №. 10. - S. 567-576.
4. Ermatov N., Okhunova M., Kamilova A. Hygienic analysis of the health of schoolchildren // world science: problems and innovations. – 2022. - S. 245-246.
5. Gavryushin M.Yu. Anthropometric features of physical development of schoolchildren in a modern megalopolis [tekst] / m.Yu. Gavryushin, I.I. Berezin, O.V.Sazonova // Kazan Medical Journal. – 2016. - T. 97. – № 4. - S.629-633.
6. Grechkina L.I. Morphofunctional features of physical development of school-age children, manifested in various climatogeographic zones of the Magadan region [tekst] / I.I. Grechkina, V.O. Karandasheva // Bulletin of the Northeastern Scientific

7. Kamilova, Aida Sheralievna, Ermatov, Nizam Jumakulovich, Kamilov, Jamshid Yuldashevich / The role of the diet of nutrition in helminthiasis, which occurs in children // ORIENSS. 2023. №4-2.
8. Ortikova M.M. The algorithm for attracting gelminthic companies from among Russian citizens to the Republic of Tajikistan / M.M. Kotikova, A. A. Akhmedov, Z.A. Mirzoeva, D.D. Davlyatova / / Bulletin of postgraduate education in the field of healthcare. G. Dushanbe. 17 November 2017. - S. 45-48.
9. Ortiqov B. B. ISHCHILARDA UCHRAYDIGAN KASALLANISHLARNING GIGIYENIK TAHLILI //Медицинский журнал молодых ученых. – 2025. – №. 16 (12). – С. 155-159.
10. Ortiqov B. B. Kaliy ishlab chiqarish korxonalarida ishlovchi ishchilarning ovqatlanish tartibiga qo ‘yiladigan gigiyenik talablar //Медицинский журнал молодых ученых. – 2025. – №. 13 (03). – С. 194-200.
11. Ortiqov B. B., Baxtiyorova G. R., Tugilova S. N. Analysis of the main risk group products in the daily diet of tekstil enterprises'employees. – 2024.
12. Shaykhova, G. I., et al. Factors that form disorders in the physiological bends of the spine of children and adolescents.” (2022).
13. Хайдаров Ш.М., Алимухамедов Д.Ш. (2025). Тез-тез касалланувчи болаларнинг кунлик рационини микронутриентлар билан бойитилгандан кейин таомноманинг самарадорлигини гигиеник баҳолаш. Медицинский журнал молодых ученых, 1(16 (12), 282–286. извлечено от <https://journals.tnmu.uz/yotj/article/view/3135>.